







### **Ulcus periwound maceration**

### After 13 Days

#### Men, 77 years of age

VG: Papillair urotheelcelca. and adenoca. Laparotomie and later on a jejuno-ileostomie and transversotransversostomie. Ending with a sufficient Abdominal wall with a granulating wound. Maceration peri wound. Insufficient Kidneyfunction.

Picture 1: Day 0 Start using Aldanex. Every time When a hyrofiber is changed.

Picture 2: Day 3 Less maceration.

Picture 3: Day 13 Ulcus closed.

Case: Groene Hart Ziekenhuis/ Stella Amesz (NP) en Annelies van Zandbergen (VS i.o.)– Gouda







## **Dermatitis**

**After 8 hours** 

Adult

Dermatitis leaking ileo fistula

Picture 1: Hour 0
Start using Aldanex

Picture 2: Hour 6

Picture 3: Hour 8











### **Pressure Ulcer**

### After 7 Days

Lady, bedridden
Change position and special Matrass.

First degree Pressure Ulcer thoracal spine due to static bedridden.\

Picture 1: Day 0 Redness, Erythema.

Start Aldanex.

Picture 2: Day 2

Significantly reducing redness / erythema in size

Picture 3: Day 7

Visible improvement, redness reduced to zero

Case: HOZO/ Jan van Soest (wondverpleegkundige) – Hillegom







## Intertrigo After 5 Days

Adult

Intertrigo, between the Toes

Picture 1: Day 0

Starting twice a day using Aldanex

Picture 2: Day 2

Picture 3: Day 5









### **Incontinence Dermatitis**

**After 4 Days** 

Patient

Baby, 9 months

Diper dermatitis

Picture 1: Day 0

Starting twice a day using Aldanex

Picture 2: Day 2

Picture 3: Day 4









### **Incontinence Dermatitis**

## After 9 days

#### **Patient**

Extreme premature, Born after 25 weeks Pregnancy

**Diper dermatitis** 

Picture 1: Day 0

Starting twice a day using Aldanex

Picture 2: Day 2

Picture 3: Day 9

